



NEXT-LEVEL PARASITE PROTECTION

Everyday life leaves dogs open to parasite risk. NexGuard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) is designed to be an easy-to-give, tasty, beef-flavored chew that offers next-level protection against fleas, ticks, heartworm disease, roundworms and hookworms.



FLEAS



TICKS



HEARTWORM DISEASE



ROUNDWORMS



HOOKWORMS



Triple-Action Formula

Afoxolaner, moxidectin, and pyrantel protect against common internal and external parasites in dogs, including the aggressive and invasive tick species, *H. longicornis*.



Gentle on Puppies

Safe for puppies at 8 weeks, weighing just 4 pounds.



Designed with Dogs in Mind

- Tasty, beef-flavored soft chew is designed to make monthly compliance easy and enjoyable.
- In a laboratory study, dogs preferred NexGuard® PLUS over Simparica Trio®.^{1*}



Safe and Effective

- In a study of MDR1-deficient Collies, no signs of avermectin toxicity were seen.²
- In a separate study, NexGuard PLUS chews were well-tolerated in heartworm-positive dogs.²

MOXIDECTIN AGAINST RESISTANT HEARTWORMS

Resistance to all macrocyclic lactones has been confirmed in rare isolates of *Dirofilaria immitis* in the United States. Studies have shown that macrocyclic lactones are more effective against resistant heartworm isolates when administered with repeated monthly doses.

In a recent study evaluating the efficacy against the resistant JYD-34 heartworm isolate, NexGuard PLUS and Simparica Trio, provided comparable levels of control (>99% efficacy).³



NEW
STUDY

against resistant
heartworm
isolates

IMPORTANT SAFETY INFORMATION: NexGuard® PLUS is for use in dogs only. The most frequently reported adverse reactions include diarrhea, vomiting, lethargy, and itching. Use with caution in dogs with a history of seizures or neurologic disorders. Dogs should be tested for existing heartworm infection prior to starting a preventive. For more information, see prescribing information or visit NexGuardPLUSClinic.com.

sound
byte

NexGuard® PLUS
(afoxolaner, moxidectin, and
pyrantel chewable tablets)

 **Boehringer
Ingelheim**

HEALTHY HABITS FOR PETS

Despite tick species extending their habitats across the US, we see that **93%** of dogs leave the vet clinic with less than 12 months of prescription flea and tick protection. And, although heartworm disease incidence continues to rise, **88%** of dogs leave the vet clinic with less than 12 months of prescription heartworm preventive product.⁴

With an emphasis on the importance of regular vet visits and veterinarian-prescribed parasite protection, the Healthy Habits for Pets website is designed to help you support pet owners as they develop lifelong habits to help keep their pets healthy.

[Visit HealthyHabitsForPets.com](https://HealthyHabitsForPets.com) to learn more.

* For dogs demonstrating a preference, they preferred NexGard® PLUS chews over Simparica Trio.

† See Satisfaction Guarantee For Terms and Conditions

1. Data on file at Boehringer Ingelheim.

2. Freedom of Information Summary, NADA 141-554. NexGard PLUS chews.

3. Prullage J, Frost J, DiCosto U, et al. Preventive efficacy of six monthly doses of NexGard® PLUS or Simparica Trio® against a macrocyclic lactone-resistant isolate (JVD-34) of *Dirofilaria immitis* and of a single dose of NexGard PLUS against a susceptible isolate. *Parasites & Vectors*. 2024;17(1). doi:https://doi.org/10.1186/s13071-024-06603-z

4. Data on file, IDEXX Laboratories Inc, Westbrook, Maine, USA

IMPORTANT SAFETY INFORMATION: NexGard® PLUS is for use in dogs only. The most frequently reported adverse reactions reported in clinical trials were diarrhea, vomiting, lethargy, and itching. NexGard PLUS contains afoxolaner, a member of the isoxazoline class, which has been associated with neurologic adverse reactions including tremors, ataxia, and seizures in dogs with or without a history of seizures. Use with caution in dogs with a history of seizures or neurologic disorders. The safe use of NexGard PLUS has not been evaluated in breeding, pregnant, or lactating dogs. Dogs should be tested for existing heartworm infection prior to starting a heartworm disease preventive. For heartworm disease prevention, give once monthly for at least 6 months after last exposure to mosquitoes. For more information, see the prescribing information or visit NexGardPLUSClinic.com.



NexGard® is a registered trademark of Boehringer Ingelheim Animal Health France, used under license. All other trademarks are the property of their respective owner. ©2025 Boehringer Ingelheim Animal Health USA Inc., Duluth, GA. All rights reserved. US-PET-0077-2025.

NexGard® PLUS
(afoxolaner, moxidectin, and
pyrantel chewable tablets)

MORE THAN MEDICINE

Boehringer Ingelheim offers resources to keep your patients protected, your clients satisfied and your clinic supported.



Client Education

Free resources to educate your clients about NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) and the importance of protection.



Satisfaction Guarantee

We want your clients to be satisfied with our products, which is why we offer a satisfaction guarantee.[†]

sound byte

CHECK-OFF &
ECHO NOTES

Be ready to talk to dog owners about the risk of parasites in dogs.

Question to ask dog owners in your clinic:

Are you familiar with NexGard PLUS?

If **YES**, confirm and discuss benefits.

If **NO**, explain why you recommend the product, and how easy it is to transition the dog to the product you prefer.

Coverage

Just one tasty monthly dose of NexGard PLUS protects dogs from fleas, ticks, heartworm disease, roundworms and hookworms.

Neutral Invitation

Let's review some of the features of NexGard PLUS...

It's the Customer's Decision

Discuss the risks of parasites in dogs. Reinforce the importance of safe, effective, One-And-Done protection with NexGard PLUS.

NexGard® PLUS

(afoxolaner, moxidectin, and pyrantel chewable tablets)

For oral use in dogs only.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Description:
NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) is available in five sizes of beef-flavored, soft chewables for oral administration to dogs and puppies according to their weight. Each chewable is formulated to provide minimum doses of 1.14 mg/lb (2.5 mg/kg) afoxolaner, 5.45 mcg/lb (12 mcg/kg) moxidectin, and 2.27 mg/lb (5.0 mg/kg) pyrantel (as pamoate salt).

Afoxolaner is a member of the isoxazoline family of compounds. Its chemical name is 1-Naphthalene-carboxamide,4-[5-[3-chloro-5-(trifluoromethyl)-phenyl]-4,5-dihydro-5-(trifluoromethyl)-3-isoxazoly]-N-[2-oxo-2-[(2,2,2-trifluoroethyl)amino]ethyl].

Moxidectin is a semisynthetic macrocyclic lactone derived from the actinomycete *Streptomyces cyaneogriseus noncyanogenus*. The chemical name for moxidectin is [6R,23E,25S(E)]-5-O-Demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28-epoxy-23-(methoxyimino) milbemycin B.

Pyrantel is a member of the tetrahydropyrimidine family of compounds. Its chemical name is (E)- 1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4, 4' methylenebis [3-hydroxy-2-naphthoate](1:1).

Indications:
NexGard® PLUS is indicated for the prevention of heartworm disease caused by *Dirofilaria immitis* and for the treatment and control of adult hookworm (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) and roundworm (*Toxocara canis* and *Toxascaris leonina*) infections. NexGard® PLUS kills adult fleas and is indicated for the treatment and prevention of flea infestations (*Ctenocephalides felis*) and the treatment and control of *Ixodes scapularis* (black-legged tick), *Rhipicephalus sanguineus* (brown dog tick), *Dermacentor variabilis* (American dog tick), *Amblyomma americanum* (lone star tick), and *Haemaphysalis longicornis* (longhorned tick) infestations for one month in dogs and puppies eight weeks of age and older, weighing four pounds of body weight or greater.

Dosage and Administration:
NexGard® PLUS is given orally once a month at the minimum dosage of 1.14 mg/lb (2.5 mg/kg) afoxolaner, 5.45 mcg/lb (12 mcg/kg) moxidectin, and 2.27 mg/lb (5.0 mg/kg) pyrantel (as pamoate salt).

For heartworm disease prevention, give once monthly for at least six months after last exposure to mosquitoes (see Effectiveness).

Dosing Schedule:

Body Weight (lbs.)	Afoxolaner Per Chewable (mg)	Moxidectin Per Chewable (mcg)	Pyrantel* Per Chewable (mg)	Chewables Administered
4 – 8	9.375	45	18.75	One
8.1 – 17	18.75	90	37.5	One
17.1 – 33	37.5	180	75	One
33.1 – 66	75	360	150	One
66.1 – 132	150	720	300	One
Over 132	Administer the appropriate combination of chewables			

*As pamoate salt.

NexGard® PLUS can be administered with or without food. Care should be taken to ensure that the dog consumes the complete dose and that part of the dose is not lost or refused. If a dose is missed, administer NexGard® PLUS and resume a monthly dosing schedule.

Heartworm Prevention:

NexGard® PLUS should be administered at monthly intervals year-round or, at a minimum, administration should start within one month of the dog's first seasonal exposure to mosquitoes and should continue at monthly intervals until at least six months after the dog's last exposure (see **Effectiveness**). When replacing another monthly heartworm preventive product, the first dose of NexGard® PLUS should be given within a month of the last dose of the former medication.

Flea Treatment and Prevention:

NexGard® PLUS should be administered year-round at monthly intervals or started at least one month before fleas become active. To minimize the likelihood of flea reinfestation, it is important to treat all animals within a household with an approved flea control product.

Tick Treatment and Control:

NexGard® PLUS should be administered year-round at monthly intervals or started at least one month before ticks become active.

Intestinal Nematode Treatment and Control:

NexGard® PLUS treats and controls adult hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) and roundworms (*Toxocara canis* and *Toxascaris leonina*). For the treatment of adult hookworm and roundworm infections, NexGard® PLUS should be administered as a single dose. Monthly use of NexGard® PLUS will control any subsequent infections. Dogs may be exposed to and can become infected with hookworms and roundworms throughout the year, regardless of season or climate.

Contraindications:

There are no known contraindications for the use of NexGard® PLUS.

Warnings:

Not for use in humans. Keep this and all drugs out of the reach of children. In case of accidental ingestion, contact a physician for medical advice.

Keep NexGard® PLUS in a secure location out of the reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

Precautions:

Afoxolaner, one of the ingredients in NexGard® PLUS, is a member of the isoxazoline class. This class has been associated with neurologic adverse reactions including tremors, ataxia, and seizures. Seizures have been reported in dogs receiving isoxazoline class drugs, even in dogs without a

history of seizures. Use with caution in dogs with a history of seizures or neurologic disorders.

Treatment with fewer than six monthly doses after the last exposure to mosquitoes has not been evaluated and may not provide complete heartworm prevention.

Prior to administration of NexGard® PLUS, dogs should be tested for existing heartworm infection. At the discretion of the veterinarian, infected dogs should be treated with an adjuvant to remove adult heartworms. NexGard® PLUS is not effective against adult *D. immitis*.

The safe use of NexGard® PLUS in breeding, pregnant, or lactating dogs has not been evaluated.

Adverse Reactions:

In a field safety and effectiveness study, NexGard® PLUS was administered to dogs for the prevention of heartworm disease. The study included a total of 272 dogs (134 administered NexGard® PLUS and 138 administered active control) treated once monthly for 11 treatments. Over the 330-day study period, all observations of potential adverse reactions were recorded. The most frequent reactions reported in the NexGard® PLUS group are presented in the following table.

Table 1: Dogs With Adverse Reactions.

Clinical Sign	NexGard® PLUS n = 134 Number (Percentage)	Active Control n = 138 Number (Percentage)
Diarrhea	9 (6.7%)	7 (5.1%)
Vomiting	6 (4.5%)	7 (5.1%)
Lethargy	3 (2.2%)	5 (3.6%)
Itching	3 (2.2%)	3 (2.2%)
Dermatitis	2 (1.5%)	1 (0.7%)
Anorexia	1 (0.7%)	4 (2.9%)
Muscle tremor	1 (0.7%)	1 (0.7%)

One dog in the NexGard® PLUS group was reported to exhibit muscle tremors along with nausea and depression for one day after the Day 0 treatment. The dog remained in the study and muscle tremors were not reported after any subsequent treatments.

Contact Information:

For a copy of the Safety Data Sheet (SDS) or to report suspected adverse drug events, contact Boehringer Ingelheim Animal Health USA Inc. at 1-888-637-4251 or www.nexgardforpets.com.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae.

Clinical Pharmacology:

Mode of Action:

NexGard® PLUS (afoxolaner, moxidectin, and pyrantel chewable tablets) contains the three active pharmaceutical ingredients afoxolaner, moxidectin, and pyrantel (as pamoate salt).

Afoxolaner is a member of the isoxazoline family, shown to bind at a binding site to inhibit insect and acarine ligand-gated chloride channels, in particular those gated by the neurotransmitter gamma-aminobutyric acid (GABA), thereby blocking pre- and postsynaptic transfer of chloride ions across cell membranes. Prolonged afoxolaner-induced hyperexcitation results in uncontrolled activity of the central nervous system and death of insects and acarines. The selective toxicity of afoxolaner between insects and acarines and mammals may be inferred by the differential sensitivity of the insects and acarines' GABA receptors versus mammalian GABA receptors.

Moxidectin is an endectocide in the macrocyclic lactone class. Moxidectin acts by interfering with chloride channel-mediated neurotransmission in susceptible parasites, which results in paralysis and death of the parasite.

Pyrantel is a nematocide belonging to the tetrahydropyrimidine class. Pyrantel acts as a depolarizing, neuromuscular-blocking agent in susceptible parasites, causing paralysis and death or expulsion of the parasite.

Pharmacokinetics:

Following a single oral administration of a near-final formulation of NexGard® PLUS (at mean doses of 3.9 mg/kg afoxolaner, 18.8 mcg/kg moxidectin, and 7.8 mg/kg pyrantel pamoate) in fed and fasted Beagle dogs (10 to 21 months of age), afoxolaner and moxidectin were more rapidly absorbed in the fasted state with a time to maximum concentration (Tmax) of 2 to 3 hours.

The afoxolaner mean maximum plasma concentrations (Cmax) in the fed and fasted states were 1610 and 2200 ng/mL (CV=33 and 16%) and the moxidectin mean Cmax values were 11.1 and 15.5 ng/mL (CV=39 and 24%), respectively. The area under the curve (AUC) for afoxolaner and moxidectin were similar between fed and fasted states. Post-dose pyrantel plasma concentrations were quantifiable out to 24 hours.

Following six oral administrations of NexGard® PLUS at 1, 3, and 5X the maximum exposure dose of 5 mg/kg, 24 mcg/kg, and 10 mg/kg afoxolaner, moxidectin, and pyrantel pamoate, respectively, every 28 days in 8-week-old Beagle dogs, afoxolaner and moxidectin Tmax ranged from 2 to 6 hours. The observed mean Cmax and AUC at steady state in the 1X dose group were 2230 ng/mL and 19000 days*ng/mL for afoxolaner and 14.8 ng/mL and 55.2 days*ng/mL for moxidectin, respectively. Based on mean Cmin, afoxolaner and moxidectin accumulated by less than 4-fold at steady state. Afoxolaner and moxidectin exposure increased in a dose proportional manner between the 1X and 3X dose groups but was less than dose proportional in the 5X dose group.

Pyrantel pamoate is poorly absorbed into systemic circulation. Pyrantel pamoate is intended to remain in the gastrointestinal tract to allow effective concentrations to be delivered to gastrointestinal nematodes.

Effectiveness:

Heartworm Prevention:

In two well-controlled laboratory studies, NexGard® PLUS was 100% effective against induced *D. immitis* infections when administered for six consecutive months.

In a well-controlled US field study consisting of 120 dogs administered NexGard® PLUS and 124 administered an active control, no dogs treated with NexGard® PLUS tested positive for heartworm disease. All dogs treated with NexGard® PLUS were negative for *D. immitis* antigen and blood microfilariae at study completion on Day 330.

Flea Treatment and Prevention:

In a well-controlled laboratory study, NexGard® PLUS demonstrated

≥99.8% effectiveness against adult fleas 24 hours after weekly infestations for one month.

In a separate well-controlled laboratory study, afoxolaner alone began to kill fleas four hours after initial administration and demonstrated >99% effectiveness at eight hours.

In an additional well-controlled laboratory study, afoxolaner alone demonstrated 100% effectiveness against adult fleas 24 hours post-infestation for 35 days and was ≥93% effective at 12 hours post-infestation through Day 21 and on Day 35. On Day 28, afoxolaner alone was 81.1% effective 12 hours post-infestation. Dogs in both the afoxolaner-treated and control groups that were infested with fleas on Day -1 generated flea eggs at 12 and 24 hours post-treatment (0-11 eggs and 1-17 eggs in the afoxolaner-treated dogs, and 4-90 eggs and 0-118 eggs in the control dogs, at 12 and 24 hours, respectively). At subsequent evaluations post-infestation, fleas from dogs in the afoxolaner-treated group were essentially unable to produce any eggs (0-1 eggs), while fleas from dogs in the control group continued to produce eggs (1-141 eggs).

In a 90-day US field study conducted in households with existing flea infestations of varying severity, the effectiveness of afoxolaner alone against fleas on the Day 30, 60, and 90 visits compared with baseline was 98.0%, 99.7%, and 99.9%, respectively. In a second 90-day US field study, the effectiveness of afoxolaner alone against fleas on the Day 30, 60, and 90 visits compared with baseline was 97.5%, 99.7%, and 99.9%, respectively. Dogs in the second study with signs of Flea Allergy Dermatitis (FAD) showed improvement in erythema, alopecia, papules, scales, crusts, and excoriation following treatment, as a direct result of eliminating fleas.

Collectively, the data from the five studies (three laboratory and two field) demonstrate that NexGard® PLUS kills fleas before they can lay eggs, thus preventing subsequent flea infestations after the start of treatment of existing flea infestations.

Tick Treatment and Control:

In well-controlled laboratory studies, NexGard® PLUS demonstrated ≥97% effectiveness against *Amblyomma americanum* 72 hours post-infestation, for one month.

In well-controlled laboratory studies, a chewable containing afoxolaner alone, one of the active ingredients in NexGard® PLUS, demonstrated effectiveness against *Ixodes scapularis*, *Rhipicephalus sanguineus*, and *Dermacentor variabilis* at 48 hours post-infestation, and against *Haemaphysalis longicornis* at 72 hours post-infestation, for one month.

Intestinal Nematode Treatment and Control:

Elimination of adult roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum*, *Ancylostoma braziliense*, and *Uncinaria stenocephala*) was demonstrated in well-controlled laboratory studies.

Target Animal Safety:

Margin of Safety:

NexGard® PLUS was administered orally at 1, 3, and 5X the maximum exposure doses at approximately 28-day intervals for six treatments to 8-week-old Beagle puppies. Dogs in the control group were sham-dosed. There were no clinically relevant, treatment-related effects on body weights, food consumption, clinical pathology (hematology, coagulation, serum chemistry, and urinalysis), gross pathology, histopathology, organ weights, or ophthalmic examinations. Mild, self-limiting diarrhea (with and without blood) was possibly related to treatment, as there were more incidences in the NexGard® PLUS groups than the control group throughout the study, including within 48 hours after treatment.

Avermectin-Sensitive Collie Safety:

NexGard® PLUS was administered orally at 1, 3, and 5X the maximum label dose to MDRI-deficient Collies once on Day 0, with a second administration to the 1X group on Day 28. Dogs in the control group were sham-dosed on Days 0 and 28. No clinical signs of avermectin toxicity were noted in any dog at any time during the study. Vomiting was observed in some dogs in the 3X and 5X groups and resolved without treatment. Diarrhea, with or without blood, was observed in some dogs in all of the NexGard® PLUS groups and resolved without treatment.

Heartworm-Positive Safety:

NexGard® PLUS was administered orally at 1X and 3X the maximum exposure doses at approximately 28-day intervals for three treatments to Beagle dogs with adult heartworm infections and circulating microfilariae. Dogs in the control group were sham-dosed. Diarrhea was observed in one dog in the 1X group and in three dogs in the 3X group, and vomiting was observed in two dogs in the 3X group. No signs of avermectin toxicity were observed at any time during the study. There were no clinical signs associated with death of the microfilariae observed in any of the dogs.

Field Safety:

In a well-controlled field study, NexGard® PLUS was used concurrently with other medications such as vaccines, antibiotics, non-steroidal anti-inflammatory drugs (NSAIDs), anesthetics, sedatives, analgesics, steroids, anthelmintics, antiemetics, and antipruritics. No adverse reactions were associated with the concurrent use of NexGard® PLUS and other medications.

How Supplied:

NexGard® PLUS is available in five strengths of beef-flavored soft chewables formulated according to the weight of the dog (see **Dosage and Administration**). Each chewable size is available in color-coded packages of 1, 3, or 6 chewables.

Storage Information:

Store in original package at or below 25°C (77°F) with excursions permitted up to 40°C (104°F).

Approved by FDA under NADA # 141-554

Marketed by: Boehringer Ingelheim Animal Health USA Inc., Duluth, GA 30096

NEXGARD® is a registered trademark of Boehringer Ingelheim Animal Health France, used under license.

©2024 Boehringer Ingelheim Animal Health USA Inc. All rights reserved. US-PET-0084-2023-V4.

181531-004
Rev. 08/2024

